Abbey Pain Scale

Resident	.'s name:									
Date and	I time:				_					
Name/De	esignation of Pers	on Completing	g the Scale: _							
Last Pair	n Relief					Dos	age:			
Time of L	₋ast Pain Relief:									
Q1	Vocalization	(e.g. whimpe	g. whimpering, groaning, crying)]	
	Absent - 0	Mild - 1	Moderate - 2	Severe - 3	3					
Q2 Facial Expression (e.g. looking tense, frowning, grimacing, looking frightened) Q2									1	
	Absent - 0	Mild - 1	Moderate - 2	Severe - 3	3					
Q3	Change in Bowithdrawn)	ody Languag	e(e.g. fidgetin	g, rocking,	guarding part of body	/,	Q3			
	Absent - 0	Mild - 1	Moderate - 2	Severe - 3	3		·		_	
Q4	Behavioral C usual patter		ncreased conf	iusion, refu	sing to eat, alternatio	n in	Q4			
	Absent - 0	Mild - 1	Moderate - 2	Severe - 3	3					
Q5		Physiological Change (e.g. temperature, pulse or blood pressure outside Q5 normal limits, perspiring, flushing or pallor)								
	Absent - 0	Mild - 1	Moderate - 2	Severe - 3	3					
Q6	Physical Changes (e.g. skin tears, pressure areas, arthritis, contractures, previous injuries)						Q6			
	Absent - 0	Mild - 1	Moderate - 2	Severe - 3	3					
						Total score	e:			
				Pain S	Severity					
			Tick the	box that ma	atches the total score.					
0-2		3-7		8-13		14+				
No Pain		Mild	Mild Moderate			Severe				
				Туре	of Pain					
			Tick the	e box that m	natches pain severity.					
Mild				Moderate			Severe			
Chronic				Acute			Acute on Chronic			

Abbey, J; De Bellis, A; Piller, N; Esterman, A; Giles, L; Parker, D and Lowcay, B.Funded by the JH and JD Gunn Medical Research Foundation 1998-2002

Modified Abbey Pain Scale

	Score, date, and time						
Vocalization (e.g. whimpering, groaning, crying)							
Absent - 0 Mild - 1 Moderate - 2 Severe - 3							
Facial Expression (e.g. looking tense, frowning, grimacing, looking frightened)							
Absent - 0 Mild - 1 Moderate - 2 Severe - 3							
Change in Body Language (e.g. fidgeting, rocking, guarding part of body, withdrawn)							
Absent - 0 Mild - 1 Moderate - 2 Severe - 3							
Behavioral Change (e.g. increased confusion, refusing to eat, alternation in usual patterns)							
Absent - 0 Mild - 1 Moderate - 2 Severe - 3							
Physiological Change (e.g. temperature, pulse or blood pressure outside normal limits, perspiring, flushing or pallor)							
Absent - 0 Mild - 1 Moderate - 2 Severe - 3							
Physical Changes (e.g. skin tears, pressure areas, arthritis, contractures, previous injuries)							
Absent - 0 Mild - 1 Moderate - 2 Severe - 3							
Total Score:							
Did the pain increase (+) or decrease (-) from initial previous test? Add (+) or (-) to score.							

Modified Abbey Pain Scale

	Score, date, and time						
Vocalization (e.g. whimpering, groaning, crying)							
Absent - 0 Mild - 1 Moderate - 2 Severe - 3							
Facial Expression (e.g. looking tense, frowning, grimacing, looking frightened)							
Absent - 0 Mild - 1 Moderate - 2 Severe - 3							
Change in Body Language (e.g. fidgeting, rocking, guarding part of body, withdrawn)							
Absent - 0 Mild - 1 Moderate - 2 Severe - 3							
Behavioral Change (e.g. increased confusion, refusing to eat, alternation in usual patterns)							
Absent - 0 Mild - 1 Moderate - 2 Severe - 3							
Physiological Change (e.g. temperature, pulse or blood pressure outside normal limits, perspiring, flushing or pallor)							
Absent - 0 Mild - 1 Moderate - 2 Severe - 3							
Physical Changes (e.g. skin tears, pressure areas, arthritis, contractures, previous injuries)							
Absent - 0 Mild - 1 Moderate - 2 Severe - 3							
Total Score:							
Did the pain increase (+) or decrease (-) from initial previous test?							
Add (+) or (-) to score.							

Pain Severity Legend

0-2	3-7	8-13	14+
No Pain	Mild	Moderate	Severe

Type of Pain

Mild	Moderate	Severe
Chronic	Acute	Acute on Chronic

Signature	of	Person	Comp	leting	the	Scale	: