

Barthel Index

Patient name: _____ Patient date of birth: _____

Raters name: _____ Date and time of rating: _____

Activity	Rating	Score
Feeding	0 = Unable	
	5 = Needs help cutting, spreading butter, etc., or requires a modified diet	
	10 = Independent	
Bathing	0 = Dependent	
	5 = Independent (or in shower)	
Grooming	0 = Needs to help with personal care	
	5 = Independent face/hair/teeth/shaving (implements provided)	
Dressing	0 = Dependent	
	5 = Needs help but can do about half unaided	
	10 = Independent (including buttons, zips, laces, etc.)	
Bowels	0 = Incontinent (or needs to be given enemas)	
	5 = Occasional accident	
	10 = Continent	
Bladder	0 = Incontinent, or catheterized and unable to manage alone	
	5 = Occasional accident	
	10 = Continent	
Toilet use	0 = Dependent	
	5 = Needs some help, but can do something alone	
	10 = Independent (on and off, dressing, wiping)	

Activity	Rating	Score
Transfer (bed to chair and back)	0 = Unable, no sitting balance	
	5 = Major help (one or two people, physical), can sit	
	10 = Minor help (verbal or physical)	
	15 = Independent	
Mobility (on level surfaces)	0 = Immobile or < 50 yards	
	5 = Wheelchair independent, including corners, > 50 yards	
	10 = Walks with help of one person (verbal or physical) > 50 yards	
	15= Independent (but may use any aid; for example, stick) > 50 yards	
Stairs	0 = Unable	
	5 = Needs help (verbal, physical, carrying aid)	
	10 = Independent	
Total score: (0 – 100)		
Guidelines		
<ol style="list-style-type: none"> 1. The index should be used as a record of a patient's actions, not as a record of what a patient could do. 2. The main aim is to establish the degree of independence from any help, physical or verbal, however minor and for whatever reason. 3. The need for supervision renders the patient not independent. 4. A patient's performance should be established using the best available evidence. Asking the patient, friends/relatives, and nurses are the usual sources, but direct observation and common sense are also important. However, direct testing is not needed. 5. Usually the patient's performance over the preceding 24-48 hours is important, but occasionally longer periods will be relevant. 6. Middle categories imply that the patient supplies over 50 percent of the effort. 7. The use of aids to be independent is allowed. 		
Additional notes		