

Depression Treatment Guidelines

General Information

Date of Issue:
Issuing Authority/Physician:
Review Date:

Diagnostic Criteria

• DSM-5 Criteria Summary

- Depressed mood most of the day, nearly every day.
 - Markedly diminished interest or pleasure in all, or almost all, activities.
 - Significant weight loss when not dieting, weight gain, or decrease or increase in appetite.
 - Insomnia or hypersomnia nearly every day.
 - Psychomotor agitation or retardation.
 - Fatigue or loss of energy.
 - Feelings of worthlessness or excessive or inappropriate guilt.
 - Diminished ability to think or concentrate, or indecisiveness.
 - Recurrent thoughts of death, recurrent suicidal ideation without a specific plan, or a suicide attempt.
- **ICD-10 Codes:** F32.0 (Mild), F32.1 (Moderate), F32.2 (Severe Without Psychotic Features), F32.3 (Severe With Psychotic Features), F33.x (Major Depressive Disorder, Recurrent)

Diagnostic Methods and Tests

- **Initial Clinical Interview:** Gather comprehensive psychiatric history, symptomatology, and functional impairment.
- **Standardized Screening Tools:** PHQ-9, Hamilton Depression Rating Scale, Beck Depression Inventory.
- **Physical Examination:** To rule out physical causes contributing to depressive symptoms.
- **Laboratory Tests:** Thyroid function tests, complete blood count (CBC), vitamin D levels, etc.
- **Additional Diagnostic Considerations:** Substance use assessment, bipolar disorder screening.

Treatment Guidelines

1. Psychotherapy

- Cognitive Behavioral Therapy (CBT)
- Interpersonal Therapy (IPT)
- Problem-Solving Therapy

2. Pharmacotherapy

- SSRIs (e.g., Sertraline, Fluoxetine)
- SNRIs (e.g., Venlafaxine, Duloxetine)
- Atypical Antidepressants (e.g., Bupropion, Mirtazapine)
- Tricyclic Antidepressants (TCAs) and Monoamine Oxidase Inhibitors (MAOIs) for treatment-resistant cases

3. Adjunctive Treatments

- Lifestyle modifications (exercise, diet)
- Electroconvulsive Therapy (ECT) for severe depression resistant to medication
- Transcranial Magnetic Stimulation (TMS)

CPT Codes for Billing

- Psychotherapy: 90832 (30 min), 90834 (45 min), 90837 (60 min)
- Pharmacologic Management: 99213-99215 (Outpatient visit for medication management)
- ECT: 90870


Monitoring and Follow-Up

- **Initial Response Check:** 2-4 weeks after starting treatment.
- **Regular Follow-Up:** Every 4-6 weeks to adjust treatment as necessary.
- **Long-Term Management:** Assess for maintenance therapy to prevent relapse.

Referral Considerations

- **Specialist Referral:** For patients with psychotic features, suicidal ideation, or complex comorbid psychiatric conditions.
- **Multidisciplinary Team:** Consideration for cases requiring coordinated care between mental health specialists, primary care physicians, and other healthcare professionals.

Additional Comments

A large, empty rectangular box with a thin black border, intended for providing additional comments or feedback. The box is currently blank.