## **DSM 5 Substance Use Disorder Criteria**

| $\sim$ | : 4  |     |    |
|--------|------|-----|----|
| ( ,    | lent | nam | œ. |
| $\sim$ |      |     |    |

Chart number:

| Criteria  | Meets<br>Criteria? | Notes / Supporting<br>Information |
|---|--------------------|-----------------------------------|
| Substance often taken in larger amounts or over a longer period than intended.  | ☐ Yes☐ No          |                                   |
| 2. Persistent desire or unsuccessful efforts to cut down or control the substance use.  | ☐ Yes☐ No          |                                   |
| 3. A great deal of time is spent in activities necessary to obtain the substance, use the substance, or recover from its effects.                         | ☐ Yes☐ No          |                                   |
| 4. Craving or a strong desire to use the substance.   | ☐ Yes☐ No          |                                   |
| 5. Recurrent substance use resulting in failure to fulfill major role obligations at work, school, or home.   | ☐ Yes☐ No          |                                   |
| 6. Continued substance use despite having persistent or recurrent social or interpersonal problems caused or exacerbated by the effects of the substance. | ☐ Yes☐ No          |                                   |
| 7. Important social, occupational, or recreational activities are given up or reduced because of substance use.   | ☐ Yes              |                                   |

| 8. Recurrent substance use in situations in which it is physically hazardous.   | ☐ Yes☐ No |  |  |  |  |
|---|-----------|--|--|--|--|
| 9. Continued use despite knowledge of having a persistent or recurrent physical or psychological problem that is likely to have been caused or exacerbated by the substance.  | ☐ Yes☐ No |  |  |  |  |
| <ul> <li>10. *Tolerance, as defined by either of the following:</li> <li>a. A need for markedly increased amounts of the substance to achieve intoxication or desired effect.</li> <li>b. Markedly diminished effect with continued use of the same substance.</li> </ul> | ☐ Yes☐ No |  |  |  |  |
| <ul><li>11. *Withdrawal, as manifested by either of the following:</li><li>a. The characteristic withdrawal for the substance.</li><li>b. The same (or closely related) substance is taken to relieve or avoid withdrawal symptoms.</li></ul>                             | ☐ Yes☐ No |  |  |  |  |
| (Disclaimer: This criterion is not considered to be met for those individuals taking prescription opiates solely under medical supervision.)  |           |  |  |  |  |
| Signed:   |           |  |  |  |  |
| Date:   |           |  |  |  |  |

Reference: Criteria from American Psychiatric Association (2013). Diagnostic and Statistical Manual of Mental Disorders, Fifth Edition. Washington, DC, American Psychiatric Association page 541