














Faces Pain Scale

Wong-Baker FACES Pain Rating Scale

This scale may also be scored from 0-5.







					
0 No hurt	2 Hurts a little bit	4 Hurts a little more	6 Hurts even more	8 Hurts a whole lot	10 Hurts the worst

Faces Pain Scale by Bieri et al.

						
0 No pain	1	2	3	4	5	6 Most severe pain

Faces Pain Scale—Revised by Hicks et al.

This scale may also be scored from 0-10, at intervals of 2: 0, 2, 4, 6, 8, 10.

					
0 No pain	1	2	3	4	5 Most severe pain

References

Bieri, D., Reeve, R. A., Champion, D. G., Addicoat, L., & Ziegler, J. B. (1990). The faces pain scale for the self-assessment of the severity of pain experienced by children: Development, initial validation, and preliminary investigation for ratio scale properties. *Pain, 41*(2), 139–150. [https://doi.org/10.1016/0304-3959\(90\)90018-9](https://doi.org/10.1016/0304-3959(90)90018-9)

Hicks, C. L., von Baeyer, C. L., Spafford, P. A., van Korlaar, I., & Goodenough, B. (2001). The Faces Pain Scale – Revised: toward a common metric in pediatric pain measurement. *Pain, 93*(2), 173–183. [https://doi.org/10.1016/S0304-3959\(01\)00314-1](https://doi.org/10.1016/S0304-3959(01)00314-1)

Wong-Baker FACES foundation. (2016). *Instructions for use*. <https://wongbakerfaces.org/instructions-use/>

Pain tracker

Patient name: _____ Age: _____

Date	Time	Pain reason	Pain level	Interventions/Medication for pain

Additional notes