Fall Risk Assessment

Patient Name:

Patient Identifier:

Date of Birth:

Date of Assessment:

| Risk Factor | Level | Score |
|--|--|--------------|
| Recent Falls | None in last 12 months | 2 |
| | One or more between 3 and 12 months ago | 4 |
| | One or more in last 3 months | 6 |
| | One or more in last 3 months whilst inpatient/ resident | 8 |
| Medications | None | 1 |
| Including anti-depressants, anti- hypertensives, sedatives, anti- Parkinson's, diuretics, or hypnotics. | One | 2 |
| | Тwo | 3 |
| | More than two | 4 |
| Psychological | None | |
| Including anxiety, depression, poor cooperation, poor insight or poor judgment, particularly regarding mobility. | Mildly affected by one or more of these | 2 |
| | Moderately affected by one or more of these | 3 |
| | Severely affected by one or more of these | 4 |
| Cognitive Status Using Hodkinson Abbreviated Mental Test Score (AMTS) | AMTS 9-10 | |
| | AMTS 7-8 | 2 |
| | AMTS 5-6 | 3 |
| | AMTS 4 or less | 4 |
| Risk Score | Scoring Low: 5-11 Medium:12-15 High: 16-20 | Total /20 |

Automatic High Risk Status Factors: If any of the below high risk status factors are selected, patient fall risk status is automatically high.

| | Recent change in functional status and/or medications affecting safe mobility (or anticipated) | |
|--------------------------------|--|--|
| | Dizziness / postural hypertension | |
| Fall Risk Status (choose one): | | |

Additional Notes:

