Fibromyalgia Diagnostic Criteria Checklist

Personal Information

| Patient Name: | Patient ID/Cha | Patient ID/Chart Number: | | |
|---|------------------------------|--------------------------|--|--|
| Date of Birth (MM/DD/YYYY): | Age: | Gender: | | |
| Phone No: | Email: | Email: | | |
| Date of Evaluation: | Physician: | | | |
| 1. Widespread Pain Index (WPI) |) | | | |
| Indicate areas where the patient has fe | elt pain in the last week. C | Check all that apply. | | |
| ☐ Jaw (left side) | ☐ Upper le | eg (left) | | |
| ☐ Jaw (right side) | ☐ Upper le | eg (right) | | |
| ☐ Shoulder (left) | ☐ Lower le | eg (left) | | |
| ☐ Shoulder (right) | ☐ Lower le | eg (right) | | |
| ☐ Upper arm (left) | ☐ Chest | | | |
| ☐ Upper arm (right) | ☐ Abdome | en | | |
| ☐ Lower arm (left) | ☐ Upper b | ack | | |
| ☐ Lower arm (right) | ☐ Lower b | ack | | |
| ☐ Hip (left) | □ Neck | | | |
| ☐ Hip (right) | | | | |
| Total WPI Score:/19 | l l | | | |
| 2a. Symptom Severity Score (S | C Coows) | | | |

Fatigue

□ 0 = No problem

| 1 = Slight or mild problems; generally mild or intermittent | | | | | | | | |
|--|--|---|--|--|--|--|--|--|
| 2 = Moderate; conside | 2 = Moderate; considerable problems; often present and/or at a moderate level | | | | | | | |
| ☐ 3 = Severe: pervasive, continuous, life disturbing problems | | | | | | | | |
| Waking Unrefreshed | | | | | | | | |
| 0 = No problem | | | | | | | | |
| ☐ 1 = Slight or mild problems; generally mild or intermittent | | | | | | | | |
| ☐ 2 = Moderate; considerable problems; often present and/or at a moderate level | | | | | | | | |
| ☐ 3 = Severe: pervasive | , continuous, life disturbi | ng problems | | | | | | |
| Cognitive Symptoms | | | | | | | | |
| 0 = No problem | | | | | | | | |
| ☐ 1 = Slight or mild problems; generally mild or intermittent | | | | | | | | |
| ☐ 2 = Moderate; considerable problems; often present and/or at a moderate level | | | | | | | | |
| ☐ 3 = Severe: pervasive, continuous, life disturbing problems | | | | | | | | |
| Enter your total score for Part 2a here: | | | | | | | | |
| 2b. Symptom Severity Score (SS Score) | | | | | | | | |
| 2b. Symptom Severi | ty Score (SS Score) | | | | | | | |
| 2b. Symptom Severi | ty Score (SS Score) Depression | ☐ Wheezing | Loss of appetite | | | | | |
| Muscle painIrritable bowel | , | □ Wheezing□ Raynaud's | □ Loss of appetite□ Rash | | | | | |
| Muscle painIrritable bowel syndrome | DepressionConstipationPain in upper | | | | | | | |
| Muscle painIrritable bowel syndromeFatigue/tiredness | DepressionConstipationPain in upper abdomen | ☐ Raynaud's | Rash | | | | | |
| Muscle pain Irritable bowel syndrome Fatigue/tiredness Thinking or remembering | DepressionConstipationPain in upper abdomenNausea | Raynaud's Hives/welts | □ Rash □ Sun sensitivity | | | | | |
| Muscle painIrritable bowel syndromeFatigue/tirednessThinking or | DepressionConstipationPain in upper abdomenNauseaNervousness | Raynaud's Hives/welts Ringing in ears | RashSun sensitivityHearing difficulties | | | | | |
| Muscle pain Irritable bowel syndrome Fatigue/tiredness Thinking or remembering problem | Depression Constipation Pain in upper abdomen Nausea Nervousness Chest pain | Raynaud's Hives/welts Ringing in ears Vomiting | RashSun sensitivityHearing difficultiesEasy bruising | | | | | |
| Muscle pain Irritable bowel syndrome Fatigue/tiredness Thinking or remembering problem Muscle Weakness Headache Pain/cramps in | DepressionConstipationPain in upper abdomenNauseaNervousness | Raynaud's Hives/welts Ringing in ears Vomiting Heartburn | RashSun sensitivityHearing difficultiesEasy bruisingHair loss | | | | | |
| Muscle pain Irritable bowel syndrome Fatigue/tiredness Thinking or remembering problem Muscle Weakness Headache Pain/cramps in abdomen | Depression Constipation Pain in upper abdomen Nausea Nervousness Chest pain Blurred vision | Raynaud's Hives/welts Ringing in ears Vomiting Heartburn Oral ulcers Loss/change in | Rash Sun sensitivity Hearing difficulties Easy bruising Hair loss Frequent urination | | | | | |
| Muscle pain Irritable bowel syndrome Fatigue/tiredness Thinking or remembering problem Muscle Weakness Headache Pain/cramps in | Depression Constipation Pain in upper abdomen Nausea Nervousness Chest pain Blurred vision Fever | Raynaud's Hives/welts Ringing in ears Vomiting Heartburn Oral ulcers Loss/change in taste | Rash Sun sensitivity Hearing difficulties Easy bruising Hair loss Frequent urination Painful urination | | | | | |

Count up the number of symptoms checked above.

Scoring:

• 0 symptoms: Score of 0

• 1 to 10 symptoms: Score of 1

• 11 to 24 symptoms: Score of 2

• 25 or more symptoms: Score of 3

Enter your score for Part 2b here: _____

Combined SS Score:

Now add Part 2a (Fatigue, Waking unrefreshed, Cognitive symptoms) AND 2b scores, and enter below.

Total Symptom Severity Score (SS Score) (0 to 12): _____

Diagnostic Criteria for Fibromyalgia

A patient meets the diagnostic criteria for fibromyalgia if the following three conditions are met:

1. Widespread Pain Index (WPI) and Symptom Severity (SS) Score Criteria

- 1a. The WPI score (Part 1) is greater than or equal to 7 AND the SS score (Part 2a & b) is greater than or equal to 5.
 OR
- **1b.** The WPI score (Part 1) is from 3 to 6 **AND** the SS score (Part 2a & b) is greater than or equal to 9.

2. Duration of Symptoms

Symptoms have been present at a similar level for at least 3 months.

3. Exclusion of Other Disorders

• You do not have a disorder that would otherwise explain the pain.

Examples:

- If your WPI (Part 1) was 9 and your SS score (Parts 2a & b) was 6, then you would meet the new FM diagnostic criteria.
- If your WPI (Part 1) was 5 and your SS score (Parts 2a & b) was 7, then you would **NOT** meet the new FM diagnostic criteria.

NOTES:

References: