

Functional Independence Measure (FIM) Assessment Tool

| Patient Information | | | |
|--|-----------|---------------------|-----------|
| Name: | | | |
| Age: | | Date of Assessment: | |
| Diagnosis: | | Date of Onset: | |
| Admitting Diagnosis: | | Length of Stay: | |
| Motor Subscale: Task Score (1 - 7) | | | |
| | ADMISSION | DISCHARGE | FOLLOW-UP |
| Self-care: | | | |
| Eating | | | |
| Grooming | | | |
| Bathing | | | |
| Dressing upper body | | | |
| Dressing lower body | | | |
| Toileting | | | |
| Sphincter Control: | | | |
| Bladder Management | | | |
| Bowel Management | | | |
| Transfer: | | | |
| Bed/chair/wheelchair | | | |
| Toilet | | | |
| Tub/shower | | | |
| Locomotion: | | | |
| Walk/wheelchair | | | |
| Stairs | | | |
| Motor Subtotal Score: | | | |
| Cognitive Subscale: Task Score (1 - 7) | | | |
| | ADMISSION | DISCHARGE | FOLLOW-UP |
| Communication: | | | |
| Comprehension | | | |
| Expression | | | |

| | ADMISSION | DISCHARGE | FOLLOW-UP |
|----------------------------------|-----------|-----------|-----------|
| Social Cognition: | | | |
| Social interaction | | | |
| Problem solving | | | |
| Memory | | | |
| Cognitive Subtotal Score: | | | |
| Overall Score: | | | |

Note: A total FIM score of 126 indicates complete independence, while a score of 18 indicates total dependence. A higher score reflects greater functional independence, with a score of 84 or above indicating a high level of independence, 36-83 indicating a moderate level of independence, and 18-35 indicating a low level of independence.

| Interpretation: |
|------------------------|
| |