


Hawkins Classification Chart

The Hawkins Classification Chart is a system used to categorize talar neck fractures. These are fractures in the thinnest cross-sectional portion of the talus, just proximal to the talar head.


Type I



Type I:

The Hawkins type I refers to a nondisplaced talar neck fracture. This condition has a 0-13% avascular necrosis (AVN) risk rate.


Type II



Type II:

A Hawkins type II injury is a talar neck fracture accompanied by subtalar dislocation, with an AVN risk rate of 20-50%.


Type III



Type III:

A Hawkins type III fracture involves full displacement of the talar neck, along with dislocations of either the subtalar or ankle joint, carrying an AVN risk between 20% and 100%.

Type IV



Type IV:

A Hawkins type IV injury refers to a talar neck fracture accompanied by subtalar, tibiotalar, and talonavicular dislocation. The associated risk of AVN ranges from 70-100%.

References

Cadogan, M. (2022, January 9). *Hawkins classification*. Life in the Fast Lane. <https://litfl.com/hawkins-classification/>

Canale, S. T., & Kelly, F. B. (1978). Fractures of the neck of the talus. Long-term evaluation of seventy-one cases. *The Journal of Bone and Joint Surgery. American Volume*, 60(2), 143–156. <https://pubmed.ncbi.nlm.nih.gov/417084/>

Hawkins, L. G. (1970). Fractures of the neck of the talus. *The Journal of Bone & Joint Surgery*, 52(5), 991–1002. https://journals.lww.com/jbjsjournal/Abstract/1970/52050/Fractures_of_the_Neck_of_the_Talus.13.aspx

Weatherford, B. (2023, December 18). *Talar neck fractures*. Orthobullets. <https://www.orthobullets.com/trauma/1048/talar-neck-fractures>