

Lachman Test

Name:	DoB:	Gender:	Date:
Cell Number:	Email:		
Address:			

History of Presenting Illness

Patient experienced the following:

- ☐ Pop or tear with injury ☐ Locking of knee ☐ Knee giving way

Physical Examination

General Examination	Left Knee:			Right Knee:			Comments:
Effusion	<input type="checkbox"/> Yes	<input type="checkbox"/> No	<input type="checkbox"/> NE	<input type="checkbox"/> Yes	<input type="checkbox"/> No	<input type="checkbox"/> NE	
Erythema	<input type="checkbox"/> Yes	<input type="checkbox"/> No	<input type="checkbox"/> NE	<input type="checkbox"/> Yes	<input type="checkbox"/> No	<input type="checkbox"/> NE	
Warmth	<input type="checkbox"/> Yes	<input type="checkbox"/> No	<input type="checkbox"/> NE	<input type="checkbox"/> Yes	<input type="checkbox"/> No	<input type="checkbox"/> NE	
Range of Motion	<input type="checkbox"/> NI	<input type="checkbox"/> Abnl		<input type="checkbox"/> NI	<input type="checkbox"/> Abnl		
Strength	<input type="checkbox"/> NI	<input type="checkbox"/> Abnl		<input type="checkbox"/> NI	<input type="checkbox"/> Abnl		
Neurovascular	<input type="checkbox"/> NI	<input type="checkbox"/> Abnl		<input type="checkbox"/> NI	<input type="checkbox"/> Abnl		
Maneuvers for ACL Tear (PV+, PV-)							
Lachman (58%, 2%)	<input type="checkbox"/> NI	<input type="checkbox"/> Abnl	<input type="checkbox"/> NE	<input type="checkbox"/> NI	<input type="checkbox"/> Abnl	<input type="checkbox"/> NE	
Pivot (69%, 4%)	<input type="checkbox"/> NI	<input type="checkbox"/> Abnl	<input type="checkbox"/> NE	<input type="checkbox"/> NI	<input type="checkbox"/> Abnl	<input type="checkbox"/> NE	
Anterior Drawer (29%, 6%)	<input type="checkbox"/> NI	<input type="checkbox"/> Abnl	<input type="checkbox"/> NE	<input type="checkbox"/> NI	<input type="checkbox"/> Abnl	<input type="checkbox"/> NE	
Maneuvers for Meniscus Injury Tear (PV+, PV-)							
McMurray (66%, 5%)	<input type="checkbox"/> NI	<input type="checkbox"/> Abnl	<input type="checkbox"/> NE	<input type="checkbox"/> NI	<input type="checkbox"/> Abnl	<input type="checkbox"/> NE	

NE = Not examined; ACL = Anterior cruciate ligament; NI = Normal; Abnl = Abnormal

Note: Predictive values (PV) for each maneuver are based on a pretest probability of 10 percent. If your clinical suspicion is higher or lower than this, then the PV should be correspondingly higher or lower.

Additional Notes:

Radiographic Decision-Making

Radiograph indicated if any of the following are true:

- ☐ Age less than 12 years ☐ Age 55 years or older ☐ Tenderness at head of fibula
- ☐ Isolated tenderness of patella (i.e., no bone tenderness of knee other than patella)
- ☐ Inability to flex knee to 90° ☐ Inability to take four weight-bearing steps (regardless of limping) at the time of injury and during examination.
- ☐ **Radiograph not indicated**
- ☐ **Radiograph indicated, findings:** _____

Assessment/Plan

Working Diagnosis

- ☐ Contusion ☐ Strain ☐ ACL tear ☐ ACL tear ☐ Medical cruciate ligament tear
- ☐ Medical meniscus injury ☐ Lateral meniscus injury ☐ Other:
- ☐ Exam limited; reevaluate in days

Orders

- ☐ Magnetic resonance imaging (MRI)
- ☐ Knee immobilizer
- ☐ Cast
- ☐ Ice
- ☐ No weight bearing for days
- ☐ Refer to:
- ☐ Recheck in days

Pain Medication:

- ☐ Acetaminophen (Tylenol): mg orally,
..... time(s)/day for days; number of refills
- ☐ Nonsteroidal anti-inflammatory drug:
..... mg orally prn for days; number of refills
- ☐ Oral narcotic:
..... mg orally prn for days; number of refills
- ☐ Other:

Physician's Signature

Date:

Common Maneuvers of the Knee for Assessing Possible Ligamentous and Meniscal Damage

Anterior drawer test (Top left). Place patient supine, flex the hip to 45 degrees and the knee to 90 degrees. Sit on the dorsum of the foot, wrap your hands around the hamstrings (ensuring that these muscles are relaxed), then pull and push the proximal part of the leg, testing the movement of the tibia on the femur. Do these maneuvers in three positions of tibial rotation: neutral, 30 degrees externally rotated, and 30 degrees internally rotated. A normal test result is no more than 6 to 8 mm of laxity.

Lachman test (Top right). Place patient supine on examining table, leg at the examiner's side, slightly externally rotated and flexed (20 to 30 degrees). Stabilize the femur with one hand and apply pressure to the back of the knee with the other hand with the thumb of the hand exerting pressure placed on the joint line. A positive test result is movement of the knee with a soft or mushy end point.

Pivot test (Bottom left). Fully extend the knee, rotate the foot internally. Apply a valgus stress while progressively flexing the knee, watching and feeling for translation of the tibia on the femur.

McMurray test (Bottom right). Flex the hip and knee maximally. Apply a valgus (abduction) force to the knee while externally rotating the foot and passively extending the knee. An audible or palpable snap during extension suggests a tear of the medial meniscus. For the lateral meniscus, apply a varus (adduction) stress during internal rotation of the foot and passive extension of the knee.

Adapted with permission from Jackson JL, O'Malley PG, Kroenke K. Evaluation of acute knee pain in primary care. Ann Intern Med. 2003;139:580.