Modified Romberg Test

The Modified Romberg Test template provides clinicians with a structured assessment tool to identify subtle balance impairments. Here's a detailed outline to guide the evaluation process:

| Patient information | |
|---|---------------------------------------|
| Name: | |
| Age: | |
| Date of assessment: | |
| Clinician: | |
| Initial setup | |
| Foot positioning: | |
| Right foot forward | Left foot forward |
| Safety support available: | |
| Yes | No |
| Test instructions | |
| 1. Eyes open stage | |
| Instruct the patient to stand in a tandem s | tance with their arms by their sides. |
| Observe the patient for 30 seconds with the second | heir eyes open. |
| Result: | |
| Stable | |
| Significant swaying | |
| Loss of balance | |
| Stepping | |
| 2. Eyes closed stage | |
| Instruct the patient to maintain the same stance while closing their eyes for 30 seconds. | |
| Observe for swaying or instability due to I | oss of visual input. |
| Result: | |
| Stable | |
| Significant swaying | |
| Loss of balance | |
| Stepping | |

| Test observations | |
|---|--|
| Swaying direction: | |
| Forward | |
| Backward | |
| Side-to-side | |
| Level of instability: | |
| Mild | |
| Moderate | |
| Severe | |
| Conclusion | |
| Positive test | |
| Negative test | |
| Interpretation: | |
| | |
| | |
| Further management | |
| Further management Immediate recommendations: | |
| | |
| | |
| Immediate recommendations: | |
| Immediate recommendations: | |