

Randot Test

Patient Information	
Name:	
Date of Birth:	Gender:
Medical History:	
Allergies:	
Medications:	

Test Procedure

Step	Notes
Explain the purpose of the Randot Test to the patient.	
Ensure proper lighting conditions in the examination room.	
Provide the patient with polarized glasses.	
Position the patient at the appropriate distance from the Randot Stereotest.	
Instruct the patient to cover one eye at a time with an occluder.	

<p>Begin with the Graded Circle Test:</p> <ol style="list-style-type: none">1. Ask the patient to identify the smallest circle they can see.2. Record the circle size in seconds of arc.3. Repeat for the other eye.	
<p>Proceed to the Geometric Shapes Test:</p> <ol style="list-style-type: none">1. Present random dot stereograms containing geometric shapes.2. Ask the patient to identify each shape seen in depth.3. Record any discrepancies between the two eyes.	
<p>Evaluate the patient's responses and record any observations or comments.</p>	
<p>Discuss the results with the patient and provide appropriate recommendations.</p>	

Interpretation


Normal Range:

_____ to _____ seconds of arc.

Abnormal Findings:

Recommendations:

Follow-up

A large, empty rectangular box with a thin black border, intended for taking notes or providing follow-up information. It occupies the upper half of the page below the 'Follow-up' header.