## **Resilience Test**

Client's full name: _		Date s	submitted:	
_				
Instructions				
stress, recover from	challenges, and m	aintain a sense of purp	n your current ability to ose and optimism in da in therapeutic or coach	ily life. It is not
or disagree with it,	based on your typ		best represents how m and behaviors. There a nost useful insight.	
Use the following so	ale to rate each ite	m:		
<ul> <li>1 = Strongly dis</li> <li>2 = Disagree</li> <li>3 = Neutral</li> <li>4 = Agree</li> <li>5 = Strongly ag</li> </ul>				
1. I find that I can be	ounce back quickly	from a challenge in my	usual routine.	
□ 1	2	3	4	5
2. I believe that I ha	ve control over my	own future.		
<u> </u>	2	3	4	5
3. I am able to stay	focused and optimi	stic in difficult situations	that arise.	
□ 1	2	3	4	5
4. I view challenges	as opportunities fo	r personal or professior	nal growth.	
□ 1	2	3	4	5
5. I can adapt well to	o changes (difficult	or not) in my life/usual	routine.	
_ 1	2	3	4	5
6. I am able to main	tain a positive outlo	ook even when facing d	fficulties.	
_ 1	2	3	4	5
7. I tend to see posi	tives in difficult situ	ations.		
□ 1	2	3	4	5

8. I feel a strong sense of purpose in my life.								
□ 1	2	3	4	5				
9. I can effectively manage stress in my daily life (personally and/or professionally).								
□ 1	2	3	4	5				
10. I believe that I can learn and grow from my mistakes.								
<u> </u>	2	3	4	5				
Scoring								
After totaling the scores for all 10 items, the result will fall between 10 and 50. The following ranges may offer a general indication of how the client perceives and reports their current resilience. These interpretations should be viewed as tentative and are best used to support clinical conversations, not as standalone conclusions.  10–20: Possibly lower resilience 21–30: May reflect emerging or situational resilience								
31–40: Potentially solid resilience 41–50: Likely higher resilience								
Disclaimer: This tool is intended for exploratory and reflective purposes only. It has not undergone formal psychometric validation or reliability testing, and therefore should not be used as a diagnostic tool. Mental health professionals are encouraged to use this in conjunction with validated instruments and sound clinical judgment.								
Additional notes								