

Stroke Treatment Guidelines

Patient Information

Patient Name:

Date of Birth:

Gender: Male Female Other:

Medical Record Number:

Date of Admission:

Date of Stroke Onset:

Risk Factors:

Allergies:

Current Medications:

Emergency Contact:

Recommended Treatments

1. Early Assessment and Diagnosis:

- Rapid neurological assessment upon admission.
- CT scan performed, confirming acute ischemic stroke in the left middle cerebral artery territory.
- NIH Stroke Scale score of 12 indicates moderate stroke severity.

2. Thrombolytic Therapy (if eligible):

- Patient is eligible for tissue plasminogen activator (tPA) within the 4.5-hour treatment window.
- tPA administered intravenously per standard protocol.

3. Endovascular Treatment Options (if applicable):

- Consultation with interventional radiology for consideration of mechanical thrombectomy due to large vessel occlusion.
- Patient scheduled for emergent angiography and thrombectomy.

4. Dual Antiplatelet Therapy:

- Aspirin and clopidogrel initiated post-thrombectomy to prevent recurrent stroke.

5. Early Management and Rehabilitation:

- Patient stabilized with blood pressure control and close monitoring of neurological status.
- Occupational therapy and physical therapy initiated within 24 hours to promote functional recovery.

Diagnostic Methods and Tests

CT Scan:

MRI:

NIG Stroke Scale (NIHSS):

Laboratory Tests:

Carotid Ultrasound:

Echocardiography:

Stroke-Related ICD and / or CPT Codes:

ICD-10 Codes:

CPT Codes:

Additional Comments