## **Timed Up and Go Test**

Name of patient:	Date of test:
Name of the tester:	
Before starting the test, you're going to need the following:	
<ul> <li>Tape to make a 3-meter-long distance marker</li> <li>An armchair to be used as the starting and ending points of the</li> <li>Ruler/yardstick to measure the distance for the tape</li> <li>Stopwatch to time your elderly patient</li> </ul>	ne test
Make sure your patient is wearing the footwear that they normally	y wear. They may use a walking aid, if needed.
Instructions:	Observations:
1. The test starts when you say <b>GO</b> .	As they go about their walk, make sure you observe your patient and note down anything you notice from the following:
<ol><li>When you say GO, they have to stand up from the chair and walk towards the distance marker</li></ol>	☐ Slow pace
<ol><li>Once they reach the distance marker, they have to turn around, walk back to the chair, and sit down</li></ol>	☐ They swayed a lot
4. They only need to walk at the normal pace that they do	<ul><li>They lost their balance</li><li>They have short strides</li></ul>
Tell your patient these instructions so they know what to do.	<ul> <li>☐ They have little to no arm swing</li> <li>☐ They shuffle</li> </ul>
On the word <b>GO</b> , start the timer.	<ul><li>They steadied themselves on a wall</li><li>En bloc turning while sitting down</li></ul>
Turn off the timer once they sit down on the chair.	☐ Had a hard time using walking aid
Make sure to record the time.	If any of these apply to the session, these could be indicative of possible neurological problems that need to be evaluated.
Allow them a practice round.  Time in seconds:	

## **Additional Notes**