

Urobilinogen In Urine Test

Medical Institution Details		
Name:		
Address:		
Phone No.:	Website:	
Patient Details		
Full Name:		Patient ID:
Date of Birth:	Gender:	Contact No.:
Email Address:		
Referred by Dr./Physician:		
Test Details		
Date Sample Collected:	Date of Analysis:	
Lab Technician:	Lab ID or Location:	

Symptoms Reported by Patient	
<input type="checkbox"/> Abdominal pain	<input type="checkbox"/> Yellowing of the skin (jaundice)
<input type="checkbox"/> Dark urine	<input type="checkbox"/> Fatigue
<input type="checkbox"/> Pale stools	<input type="checkbox"/> Other:

Other Relevant Readings (if taken)		
Parameter	Result	Reference Range
Bilirubin in Urine	mg/dL	Typically Negative
Liver Function Tests		Varies based on specific test

Test Results		
Parameter	Result	Reference Range
Urobilinogen in Urine	mg/dL	Typically < 1 mg/dL (but may vary based on lab)

Interpretation	
Within Normal Range	Urobilinogen levels are within the typical range, indicating no abnormal breakdown of hemoglobin in the body.
Elevated	Increased levels of urobilinogen may suggest liver diseases, hemolytic anemia, or other conditions that cause an increased breakdown of hemoglobin.
Decreased or Absent	Low or absent urobilinogen levels might indicate a blockage in the liver or bile ducts, decreased liver function, or certain types of poisoning.

Comments

Recommendations (if any)

Disclaimer: The Urobilinogen in Urine test provides insights into liver function and the breakdown of hemoglobin in the body. It's essential to interpret the results in conjunction with clinical observations and other diagnostic evaluations. Any concerns or anomalies should be discussed with a healthcare professional.

Signature of Lab Technician:	Date:
Signature of Supervising Pathologist:	Date:

Note: This document is valid only with the signatures of the respective professionals. Patient or consultative doctor should always discuss the results for a comprehensive understanding and next steps.